

SINGULAR CURVES OVER A FINITE FIELD AND WITH MANY POINTS

E. BALLICO

ABSTRACT. Recently Fukasawa, Homma and Kim introduced and studied certain projective singular curves over $\mathbb{F}_q$ with many extremal properties. Here we extend their definition to more general non-rational curves.

1. INTRODUCTION

Fix a prime p and a p -power q . Recently S. Fukasawa, M. Homma and S. J. Kim introduced a family of singular rational curves defined over $\mathbb{F}_q$, with many singular points over $\mathbb{F}_q$ and, conjecturally, some extremal properties. In this paper we discuss a similar type of curves, discuss their extremal properties and, in some cases, show that they are, more or less, the curves introduced in [5]. The zeta-function $Z_Y(t)$ of a singular curve Y is explicitly given in terms of the Frobenius on a “topological” invariant $H_c^1(Y, \mathbb{Q}_\ell)$ ([2], [1], p. 2, [8], [12]). Hence $Z_Y(t)$ does not detect the finer invariants of the singular points of Y (it does not distinguish between unibranch points defined over the same extension of $\mathbb{F}_q$; in particular it does not distinguish between a smooth point and a cusp). Using gluing of points of the normalization with the same residue field we may define a “minimal” singular curve with prescribed normalization and prescribed zeta-function.

Let Y be a geometrically integral projective curve defined over $\mathbb{F}_q$. Let $u : C \rightarrow Y$ denote the normalization. Since any finite field is perfect, C and u are defined over $\mathbb{F}_q$. Hence for every integer $n \geq 1$ we have $u(C(\mathbb{F}_{q^n})) \subseteq Y(\mathbb{F}_{q^n})$ and for each $P \in Y(\mathbb{F}_{q^n})$ the scheme $u^{-1}(P)$ is defined over $\mathbb{F}_{q^n}$. Hence the finite set $u^{-1}(P)_{red}$ is defined over $\mathbb{F}_{q^n}$ (but of course if $\sharp(u^{-1}(P)_{red}) > 1$ the points of $u^{-1}(P)_{red}$ may only be defined over a larger extension of $\mathbb{F}_q$). We are interested in properties of the set $Y(\mathbb{F}_q)$ knowing C . A. Weil’s study of the zeta-function of smooth projective curves was extended to the case of singular curves ([2]). We will use the very useful and self-contained treatment given by Y. Aubry and M. Perret ([1]). There are infinitely many curves Y' defined over $\mathbb{F}_q$, with C as their normalization and with the same zeta-function (see Examples 1, 2 and Lemma 2). However, given Y , there is one natural such curve if we prescribe also the sets $u^{-1}(P)_{red}$ as subsets of $C(\overline{\mathbb{F}}_q)$. Let $w_s : Y_s \rightarrow Y$ be the seminormalization of Y ([2], [11]). We recall that Y_s is an integral projective curve with C as its normalization and that $u = w_s \circ u_s$, where $u_s : C \rightarrow Y_s$ is the normalization map. Over an algebraically closed field the one-dimensional seminormal singularities with embedding dimension $n \geq 1$ are exactly the singularities formally isomorphic to the local ring at the origin of the union of the coordinate axis in $\mathbb{A}^n$. Even over a finite field the curve we will study

1991 *Mathematics Subject Classification.* 14G05; 14G10; 14G15; 14H99.

Key words and phrases. singular curve; finite field; rational point.

The author was partially supported by MIUR and GNSAGA of INdAM (Italy).

is defined in this way, i.e. the curves $C_{[q,n]}$, $n \geq 2$, defined below are obtained in the same way, i.e. the gluing process introduced by C. Traverso ([9]) gives always a seminormal curve and if the base field is algebraically closed, then all seminormal curve singularities are obtained in this way (over an algebraically closed base field a more general construction is given in [7], p. 70). We call axial singularities the curve singularities obtained in this way. Hence by definition we say that (Y, P) is an axial singularity with embedding dimension n if and only if over $\overline{\mathbb{F}}_q$ it is formally isomorphic at P to the germ at 0 of the union of the n axis of $\mathbb{A}^n$. An axial singularity of embedding dimension $n > 2$ is not Gorenstein. An axial singularity of embedding dimension 2 is an ordinary double point except that over a non-algebraically closed base field, say $\mathbb{F}_q$, we need to distinguish if the two branches of Y at P (or the two lines of its tangent cone) are defined over $\mathbb{F}_q$ or not (in the latter case each of them is defined over $\mathbb{F}_{q^2}$). Similarly, for an axial singularity (Y, P) of embedding dimension $t \geq 2$ defined over $\mathbb{F}_q$, the t lines of the tangent cone $C(P, Y)$ are defined over $\mathbb{F}_{q^t}$ and their union is defined over $\mathbb{F}_q$. In the examples we are interested in, none of these lines will be defined over a field $\mathbb{F}_{q^e}$ with $e < t$. If $P \in Y$ is a singular point, then we may associate a non-negative integer $p_a(Y, P)$ (usually called the arithmetic genus of the singularity or the drop of genus the singular point P) such that $p_a(Y) = p_a(C) + \sum_{P \in \text{Sing}(Y)} p_a(Y, P)$. When Y is an axial singularity with embedding dimension n , then $p_a(Y, P) = n - 1$.

We fix q , C and an integer $n \geq 2$. We construct a singular curve $C_{[q,n]}$ with C as its normalization and $\sharp(C_{[q,n]}(\mathbb{F}_q))$ very large in the following way. Fix an integer t such that $2 \leq t \leq n$. For each $P \in C(\mathbb{F}_{q^t}) \setminus C(\mathbb{F}_{q^{t-1}})$ the orbit of the Frobenius F_q has order t , say $\{P, \dots, F_q^{t-1}(P)\}$. Let $C_{[q,n]}$ be the only curve obtained by gluing each of these orbits (for all possible $t \leq n$) (see Remark 4). By construction $C_{[q,n]}$ is a seminormal curve defined over $\mathbb{F}_q$, each singular points of $C_{[q,n]}$ is defined over $\mathbb{F}_q$ and $\sharp(C_{[q,n]}(\mathbb{F}_q)) = N_1 + \sum_{i=2}^n (N_i - N_{i-1})/i$, where $N_i := \sharp(C(\mathbb{F}_{q^i}))$. Fix $P \in C_{[q,n]}$ with embedding dimension $t \geq 2$. The Frobenius F_q acts on the local ring $\mathcal{O}_{C_{[q,n]}, P}$ and hence on the t branches of $C_{[q,n]}$ at P (i.e. the t smooth branches through 0 of the tangent cone of $C_{[q,n]}$ at P). Since P is an axial singularity, the action of Frobenius is the restriction to $u^{-1}(P)$ of the action of the Frobenius $F_q : C(\overline{\mathbb{F}}_q) \rightarrow C(\overline{\mathbb{F}}_q)$. Hence this action is cyclic, i.e. it has a unique orbit. Thus if $O = u(P)$ with $P \in C(\mathbb{F}_{q^t}) \setminus C(\mathbb{F}_{q^{t-1}})$, then $p_a(C_{[q,n]}) = t - 1$ and none of the t branches of $C_{[q,n]}$ at $u(O)$ is defined over a proper subfield of $\mathbb{F}_{q^t}$. See Propositions 2, 3, Question 1 and Remark 2 for the relations between $\mathbb{P}_{[q,n]}^1$ and the curves B and B_n studied in [5].

2. THE CURVES $C_{[q,n]}$ AND THEIR MAXIMALITY PROPERTIES

Let $u : C \rightarrow Y$ denote the normalization map. We often write $u^{-1}(P)$ instead of $u^{-1}(P)_{\text{red}}$. Set $\Delta_Y := \sharp(u^{-1}(\text{Sing}(Y)(\overline{\mathbb{F}}_q))) - \sharp(\text{Sing}(Y)(\overline{\mathbb{F}}_q))$. The zeta-function $Z_Y(t)$ of Y is the product of the zeta-function $Z_C(t)$ of C and a degree Δ_Y polynomials whose inverse roots are roots of unity ([2], [1], Theorem 2.1 and Corollary 2.4). Let ω_i , $1 \leq i \leq 2g$, be the inverse roots of numerator of $Z_C(t)$ and β_j , $1 \leq j \leq \Delta_Y$ the inverse roots of the polynomial $Z_Y(t)/Z_C(t)$. For every integer $n \geq 1$ we have

$$(1) \quad \sharp(Y(\mathbb{F}_{q^n})) = q^n + 1 - \sum_{i=1}^{2g} \omega_i^n - \sum_{j=1}^{\Delta_Y} \beta_j^n$$

We have $\sharp(C(\mathbb{F}_{q^n})) = q^n + 1 - \sum_{i=1}^{2g} \omega_i^n$. Recall that $|\beta_j| = 1$ for all j . Hence among all curves with fixed normalization C and with fixed Δ_Y the integer $\sharp(Y(\mathbb{F}_q))$ is maximal (resp. minimal) for a curve with $\beta_j = -1$ for all j (resp. $\beta_j = 1$) for all j , if any such curve exists. If $\beta_j = -1$ for all j , then for all odd (resp. even) positive integers n among all curves with fixed normalization C and with fixed Δ_Y the value $\sharp(Y(\mathbb{F}_{q^n}))$ is maximal (resp. minimal). If $\beta_j = 1$ for all j , then for every $n > 0$ the value $\sharp(Y(\mathbb{F}_{q^n}))$ is minimal among all curves with fixed normalization C and with fixed Δ_Y .

Lemma 1. *Let Y be a geometrically integral projective curve and $u : C \rightarrow Y$ its normalization. The degree Δ_Y polynomial $Z_Y(t)/Z_C(t)$ has all its inverse roots equal to -1 if and only if for each $P \in \text{Sing}(Y)$ either $\sharp(u^{-1}(P)) = 1$ or $P \in Y(\mathbb{F}_q)$ and $u^{-1}(P)$ is formed by two points of $C(\mathbb{F}_{q^2})$ (in the latter case these two points are exchanged by the Frobenius and they are in $C(\mathbb{F}_{q^2}) \setminus C(\mathbb{F}_q)$).*

Proof. For each $Q \in C(\overline{\mathbb{F}}_q)$ let d_Q be the minimal integer $x > 0$ such that $Q \in C(\mathbb{F}_{q^x})$. The explicit form of the polynomial $Z_Y(t)/Z_C(t)$ is given in [1], Theorem 2.1. $Z_Y(t)/Z_C(t)$ is a product of polynomials, each of them associated to a different singular point of Y . Hence it is sufficient to consider separately the contribution of each singular point of Y . Fix $P \in \text{Sing}(Y)$ and call $Z_P(t)$ the associated polynomial. Let d_P be the minimal integer $t \geq 1$ such that $P \in Y(\mathbb{F}_{q^t})$. We have $(1 - t^{d_P})Z_P(t) = \prod_{Q \in u^{-1}(P)} (1 - t^{d_Q})$. Since Y is defined over $\mathbb{F}_q$, the orbit of P by the Frobenius of Y has order d_P . For any point $P' \neq P$ in this orbit, say $F_q^x(P)$ for some $x \in \{1, \dots, d_P - 1\}$ we have $u^{-1}(P') = F_q^x(u^{-1}(P))$. Hence $d_Q \geq d_P$ for each $Q \in u^{-1}(P)$ and for each $Q \in u^{-1}(P')$.

First assume $\sharp(u^{-1}(P)) = 1$. The only point, Q , of $u^{-1}(P)$ is defined over $\mathbb{F}_{q^{d_P}}$. Since $d_{u(Q)} \leq d_Q$, we get $d_Q = d_P$. We easily get that $\sharp(u^{-1}(P)) = 1$ if and only if the constant 1 is the factor of $Z_Y(t)/Z_C(t)$ associated to the orbit of P . Hence from now on we assume $\alpha := \sharp(u^{-1}(P)) \geq 2$.

If $d_Q > d_P$ for some $Q \in u^{-1}(P)$ and either $d_P \geq 2$ or $d_Q \geq 3$, then we get that $(1 - t^{d_P})Z_P(t)$ has a root of order $> \max\{d_P, 2\}$ and hence $Z_P(t)$ has a root $\neq -1$.

Now assume $d_P \geq 2$ and $d_Q = d_P$ for all $Q \in u^{-1}(P)$. We get $Z_P(t) = (1 - t^{d_P})^\alpha$. Since we assumed $\alpha \geq 2$, even in this case $Z_P(t)$ has a root $\neq -1$.

Now assume $d_P = 1$. It remains to analyze the case $d_Q \in \{1, 2\}$ for any $Q \in u^{-1}(P)$. If $d_Q = 1$ for at least one $Q \in u^{-1}(P)$, then $Z_P(1) = 0$. If $d_Q = 2$ for all $Q \in u^{-1}(P)$, then $Z_P = (1 + t)(1 - t^2)^{\alpha-1}$ has $\alpha - 1$ roots equal to 1. $\square$

Remark 1. Fix q, g, C with genus g and an integer $n \geq 2$. Set $N_i := \sharp(C(\mathbb{F}_{q^i}))$. We have $\sharp(C_{[q,n]}(\mathbb{F}_q)) = N_1 + \sum_{i=2}^n (N_i - N_{i-1})/i$, i.e.

$$(2) \quad \sharp(C_{[q,n]}(\mathbb{F}_q)) = \sum_{i=1}^{n-1} \frac{N_i}{(i+1)} + N_n/n$$

Now assume that $g > 0$ and that q is a square. If C is a minimal curve for $\mathbb{F}_q$, then it is a minimal curve for each $\mathbb{F}_{q^i}$, $i \geq 2$ (use that C is minimal if and only if $Z_C(t) = \frac{(t-q)^{2g}}{(1-t)(1-qt)}$ ([10])). Hence for fixed q and n the integer $\sharp(C_{[q,n]}(\mathbb{F}_q))$ is minimal varying C among all smooth curves of genus g if and only if C is a minimal curve. Now we assume $n \geq 3$ and generalize the construction of the curve $C_{[q,n]}$. Fix a positive integer $s \leq n - 2$ and integers $t_1, \dots, t_s$ such that $2 \leq t_1 < \dots < t_s \leq n$.

Let $C_{[q;t_1,\dots,t_s]}$ be the curve obtained from C gluing only the F_q -orbits of the points of $C(\mathbb{F}_{q^t}) \setminus C(\mathbb{F}_{q^{t-1}})$. We get

$$\#(C_{[q;t_1,\dots,t_s]}(\mathbb{F}_q)) = N_1 + \sum_{t \in \{t_1,\dots,t_s\}} (N_t - N_{t-1})/t.$$

Recall that N_1 is maximal if and only if each N_t with t odd is maximal and each N_i with i even is minimal ([10], [5], Theorem 10.1). Hence if all t_i are odd, then $\#(C_{[q;t_1,\dots,t_s]})$ is maximal if and only if C is a maximal curve.

In the case $g = 0$ we get the following result.

Proposition 1. *For every integer $n \geq 2$ we have $\#(\mathbb{P}_{[q,n]}^1) = q+1 + \sum_{t=2}^n (q^t - q^{t-1})/t$ and $p_a(\mathbb{P}_{[q,n]}^1) = \sum_{t=2}^n (q^t - q^{t-1})(t-1)/t$.*

Proposition 2. *The curve $\mathbb{P}_{[q,2]}^1$ is isomorphic over $\mathbb{F}_q$ to the plane curve $B \subset \mathbb{P}^2$ defined in [4] and [5].*

Proof. Let $u : \mathbb{P}^1 \rightarrow \mathbb{P}_{[q,2]}^1$ denote the normalization map. The normalization map $\Phi : \mathbb{P}^1 \rightarrow B$ is unramified, because the composition of it with the inclusion $B \hookrightarrow \mathbb{P}^2$ is unramified (part (i) of [5], Theorem 2.2). By [5], part (iii) of Theorem 2.2, B is a degree $q+1$ plane curve with $(q^2 - q)/2$ singular points and $\Phi(P) = \Phi(Q)$ with $P \neq Q$ if and only if $u(P) = u(Q)$. Hence the universal property of the seminormalization gives the existence of a morphism $\psi : \mathbb{P}_{[q,2]}^1 \rightarrow B$ such that ψ is a bijection. Since $p_a(\mathbb{P}_{[q,2]}^1) = (q^2 - q)/2$, we have $p_a(\mathbb{P}_{[q,2]}^1) = p_a(B)$. Hence ψ is an isomorphism. $\square$

Proposition 3. *Fix an integer $n \geq 3$. Then $\mathbb{P}_{[q,n]}^1$ is the seminormalization of the curve $B_n \subset \mathbb{P}^n$, defined in [5], §6, and there is a birational morphism $\psi_{q,n} : \mathbb{P}_{[q,n]}^1 \rightarrow B_n$ defined over $\mathbb{F}_q$ such that $\psi_{n,q} : \mathbb{P}_{[q,n]}^1(K) \rightarrow B_n(K)$ is bijective for every field $K \supseteq \mathbb{F}_q$.*

Proof. Let $u : \mathbb{P}^1 \rightarrow \mathbb{P}_{[q,n]}^1$ and $\Phi_n : \mathbb{P}^1 \rightarrow B_n$ denote the normalization maps. By [5], part (ii) of Theorem 6.4, each point $P \in \text{Sing}(B_n)$ corresponds to an integer $t \in \{2, \dots, n\}$ and an orbit of the Frobenius on $\mathbb{P}^1(\mathbb{F}_{q^t}) \setminus \mathbb{P}^1(\mathbb{F}_{q^{t-1}})$. Hence the definition of $\mathbb{P}_{[q,n]}^1$ and the universal property of the seminormalization gives a birational morphism $\psi_{q,n} : \mathbb{P}_{[q,n]}^1 \rightarrow B_n$ defined over $\mathbb{F}_q$ such that $\psi_{n,q} : \mathbb{P}_{[q,n]}^1(K) \rightarrow B_n(K)$ is bijective for every field $K \supseteq \mathbb{F}_q$. $\square$

Question 1. We guess that $\psi_{q,n}$ is an isomorphism.

Remark 2. Fix a prime power q and the integer $n \geq 3$. Let $\Phi_n : \mathbb{P}^1 \rightarrow B_n$ denote the normalization map. By [5], part (i) of Theorem 6.4, Φ_n is unramified (this is a necessary condition for being $\psi_{q,n}$ an isomorphism). The following conditions are equivalent:

- (i) the morphism $\psi_{q,n}$ is an isomorphism;
- (ii) $p_a(B_n) = p_a(\mathbb{P}_{[q,n]}^1)$;
- (iii) for each $P \in \text{Sing}(B_n)$, say with $P = \Phi_n(Q)$ and $Q \in \mathbb{P}^1(\mathbb{F}_{q^t}) \setminus \mathbb{P}^1(\mathbb{F}_{q^{t-1}})$, the singularity (B_n, P) has arithmetic genus $t-1$;
- (iv) for each $P \in \text{Sing}(B_n)$, say with $P = \Phi_n(Q)$ and $Q \in \mathbb{P}^1(\mathbb{F}_{q^t}) \setminus \mathbb{P}^1(\mathbb{F}_{q^{t-1}})$ the tangent cone $C(P, B_n) \subset \mathbb{P}^n$ is formed by t lines through P spanning a t -dimensional linear subspace.

Part (iv) is just the definition of seminormal singularity given in [2]. Since Φ_n is unramified, B_n has at P t smooth branches.

Proposition 4. *Let C be a smooth and geometrically irreducible projective curve defined over $\mathbb{F}_q$. Set $2\delta := \sharp(C(\mathbb{F}_{q^2})) - \sharp(C(\mathbb{F}_q))$. Let Y a projective curve defined over $\mathbb{F}_q$ with C as its normalization. We have $\sharp(Y(\mathbb{F}_q)) \geq \sharp(C(\mathbb{F}_q)) + \delta$ and $p_a(Y) \leq g + \delta$ if and only if Y is isomorphic to $C_{[q,2]}$ over $\mathbb{F}_q$.*

Proof. The “if” part is true, because $p_a(C_{[q,2]}) = g + \delta$ and $\sharp(C_{[q,2]}(\mathbb{F}_q)) = g + \delta$. Assume $\sharp(Y(\mathbb{F}_q)) \geq \sharp(C(\mathbb{F}_q)) + \delta$ and $p_a(Y) \leq g + \delta$. Let $u : C \rightarrow Y$ be the normalization map. The morphism u is defined over $\mathbb{F}_q$, i.e. over a field on which Y is defined, because any finite field is perfect. We have $\sharp(\text{Sing}(Y)) \leq p_a(Y) - \delta$ and equality holds only if each singular point of Y is formally isomorphic over $\overline{\mathbb{F}_q}$ either to a node or an ordinary cusp. Set $\Delta_Y := \sharp(u^{-1}(\text{Sing}(Y)(\overline{\mathbb{F}_q})) - \sharp(\text{Sing}(Y)(\overline{\mathbb{F}_q}))$. The polynomial $Z_Y(t)/Z_C(t)$ has degree Δ_Y and $\sharp(Y(\mathbb{F}_q)) \leq \sharp(C(\mathbb{F}_q)) + \Delta_Y$ and equality holds if and only if each inverse root of $Z_Y(t)/Z_C(t)$ is equal to -1 . Since $\Delta_Y \leq p_a(Y) - g$, we get $\Delta_Y = \delta$ and $p_a(Y) = g + \delta$. Since $p_a(Y) = g + \Delta_Y$ and $\sharp(\text{Sing}(Y)(\mathbb{F}_q)) \geq \delta$, we get $\text{Sing}(Y)(\overline{\mathbb{F}_q}) = \text{Sing}(Y)(\mathbb{F}_q)$, $\sharp(\text{Sing}(Y)(\mathbb{F}_q)) = \delta$ and that for each $P \in \text{Sing}(Y)(\mathbb{F}_q)$ the set $u^{-1}(P)$ is formed by two points of $C(\mathbb{F}_{q^2}) \setminus C(\mathbb{F}_q)$ exchanged by the Frobenius (Lemma 1). Since $p_a(Y) = g + \sharp(\text{Sing}(Y)(\overline{\mathbb{F}_q}))$ and $\sharp(u^{-1}(P)) \geq 2$ for each $P \in \text{Sing}(Y)(\overline{\mathbb{F}_q})$, we also get that Y is nodal. Hence Y is seminormal. The structure of the fibers of $u^{-1}(P)$, $P \in \text{Sing}(Y)(\overline{\mathbb{F}_q})$, gives $Y = C_{[q,2]}$. $\square$

Proposition 5. *Let Y be a geometrical integral projective curve defined over $\mathbb{F}_q$ and with only seminormal singularity. Let $u : C \rightarrow Y$ be the normalization map. Let Δ_Y be the degree of the polynomial $Z_Y(t)/Z_C(t)$. Assume $2\Delta_Y \leq \sharp(C(\mathbb{F}_{q^2})) - \sharp(C(\mathbb{F}_q))$. We have $\sharp(Y(\mathbb{F}_q)) \leq \sharp(C(\mathbb{F}_{[q,2]}))$ and equality holds if and only if Y is isomorphic to $C_{[q,2]}$ over $\mathbb{F}_q$.*

Proof. We have $\sharp(Y(\mathbb{F}_q)) \leq \sharp(C(\mathbb{F}_q)) + \Delta_Y$ and equality holds if and only if each inverse root of $Z_Y(t)/Z_C(t)$ is equal to -1 . Hence we may assume $2\Delta_Y = \sharp(C(\mathbb{F}_{q^2})) - \sharp(C(\mathbb{F}_q))$. Since Y has only seminormal singularities, u is unramified. Since u is unramified, we have $\sharp(u^{-1}(P)) \geq 2$ for all $P \in \text{Sing}(Y)$. Hence Lemma 1 gives that the fibers of u are the fibers of the normalization map $C \rightarrow C_{[q,2]}$. Since Y and $C_{[q,2]}$ are seminormal, we get that they are isomorphic. They are isomorphic over $\mathbb{F}_q$, because u is defined over $\mathbb{F}_q$ and the seminormalization is defined over $\mathbb{F}_q$. $\square$

Remark 3. In the case $C \cong \mathbb{P}^1$ Propositions 4 and 5 are partial answers to a questions raised in [5], Remark 2.5. Examples 1, 2 and Lemma 2 show that we need to add some conditions on the curve Y , not only to fix the normalization $\mathbb{P}^1$ and assume $\Delta_Y \leq (q^2 - q)/2$.

Example 1. Fix a geometrically integral projective curve A defined over $\mathbb{F}_q$ and $P \in A(\mathbb{F}_q)$. Now we define a geometrically integral curve Y defined over $\mathbb{F}_q$ and a morphism $v : A \rightarrow Y$ defined over $\mathbb{F}_q$, such $v|_{A \setminus \{P\}}$ is an isomorphism onto $Y \setminus v(P)$, but v is not a isomorphism. Notice that for each such pair (Y, v) we would have $p_a(Y) > p_a(A)$ and that for every integer $t \geq 1$ v induces a bijection $A(\mathbb{F}_{q^t}) \rightarrow Y(\mathbb{F}_{q^t})$. To define Y and v it is sufficient to define them in a neighborhood of P in A and the glue to it the identity map $A \setminus \{P\} \rightarrow A \setminus \{P\}$. Fix an embedding $j : A \hookrightarrow \mathbb{P}^r$, $r \geq 3$, and take a projection of $j(A)$ into $\mathbb{P}^2$ from an $(r-3)$ -dimensional

linear subspace not containing $j(P)$, but intersecting the Zariski tangent space of $j(A)$ at $j(P)$.

Example 2. Fix a geometrically integral projective curve A defined over $\mathbb{F}_q$ and any point $P \in A_{\text{reg}}(\overline{\mathbb{F}}_q)$. Let t be the minimal integer $t \geq 1$ such that $P \in A(\mathbb{F}_{q^t})$. We assume $t \geq 2$, because the case $t = 1$ is covered by Example 1. Hence the orbit of P by the action of the Frobenius F_q has order t (it is $\{P, F_q(P), \dots, F_q^{t-1}(P)\}$). Let Y denote the only curve and $v : A \rightarrow Y$ the only morphism obtained in the following way. We fix a bijection of sets $v : A \rightarrow Y$ and use it to define a topology on the set Y . Now we define Y as a ringed space. On $Y \setminus v(\{P, F_q(P), \dots, F_q^{t-1}(P)\})$ we assume that v is an isomorphism of local ringed spaces. For each $Q \in \{P, F_q(P), \dots, F_q^{t-1}(P)\}$ we impose that $\mathcal{O}_{Y,Q}$ is the local ring of a unibranch singular point and that v is the normalization map (it may be done using the method of Example 1 with Q instead of P and $\mathbb{F}_{q^t}$ instead of $\mathbb{F}_q$). We need to do the construction simultaneously over all $Q \in \{P, F_q(P), \dots, F_q^{t-1}(P)\}$ and in such a way that the morphism is defined over $\mathbb{F}_q$. As in Example 1 it is sufficient to define $v|_U$, where U is a neighborhood of $\{P, F_q(P), \dots, F_q^{t-1}(P)\}$. There is an embedding $j : A \rightarrow \mathbb{P}^r$, $r \geq t + 2$, such that the t lines $T_Q(j(A)$, $Q \in \{P, F_q(P), \dots, F_q^{t-1}(P)\}$, are linearly independent. Since j is defined over $\mathbb{F}_q$ the Frobenius F_q of $\mathbb{P}^r$ acts on $j(A)$ and on the tangent developable of A . Since $j(P)$ is defined over $\mathbb{F}_{q^t}$. Hence $T_{j(P)}j(A)(\mathbb{F}_{q^t}) \setminus j(P)$ has $(q^{t+1} - 1)/(q - 1) - 1$ elements. Fix any $O \in T_{j(P)}j(A)(\mathbb{F}_{q^t}) \setminus j(P)$. For each $x \in \{1, \dots, t - 1\}$, $F_q^x(O) \in T_{j(F_q^x(P))}j(A)(\mathbb{F}_{q^t}) \setminus j(F_q^x(P))$. Since the t tangent lines are linearly independent, the linear space $E := \langle \{O, F_q(O), \dots, F_q^{t-1}(O)\} \rangle$ has dimension $t - 1$. Since E is F_q -invariant, it is defined over $\mathbb{F}_q$. Let $\pi : \mathbb{P}^r \setminus E \rightarrow \mathbb{P}^{r-t}$ denote the linear projection from E . Since E is defined over $\mathbb{F}_q$, π is defined over $\mathbb{F}_q$. Hence the integral projective curve $T := \overline{\pi(j(A) \setminus E \cap j(A))} \subset \mathbb{P}^{r-t}$ is defined over $\mathbb{F}_q$. Since the t tangent lines are linearly independent and $O \neq j(P)$, we have $E \cap \{j(P), \dots, j(F_q^{t-1}(P))\} = \emptyset$. Hence $E \cap j(U) = \emptyset$ for a sufficiently small neighborhood U of $\{P, F_q(P), \dots, F_q^{t-1}(P)\}$. Assume for the moment that $\pi|_{j(A) \setminus j(A) \cap E}$ is birational onto its image. Since $\pi|_{j(A) \setminus j(A) \cap E}$ is birational onto its image, it is separable. Hence only finitely many points of $j(A_{\text{reg}})$ have a tangent line intersecting E . Restricting if necessary $U \subseteq A_{\text{reg}}$ we may assume that for no other point $Q \in j(U)(\overline{\mathbb{F}}_q)$ the Zariski tangent space $T_{j(Q)}(j(A))$ intersects E . Since $\pi|_{j(A) \setminus j(A) \cap E}$ is birational onto its image, it is generically injective. Hence restricting $U \subseteq A_{\text{reg}}$ we may assume that $\pi|_{j(U)}$ is injective and an isomorphism outside $\{j(P), j(F_q(P)), \dots, j(F_q^{t-1}(P))\}$. At these points the curve T has a cusp, but perhaps not an ordinary cusp, i.e. it is a unibranch singular point. Hence to conclude the example it is sufficient to find j such that $\pi|_{j(A) \setminus j(A) \cap E}$ is birational onto its image. We take as j is a linearly normal embedding of degree $d > \max\{2p_a(A) - 2, p_a(A) + t\}$. Since $d > \max\{2p_a(A) - 2, p_a(A) + t\}$, Riemann-Roch gives $r = d - p_a(A)$. Assume that $\pi|_{j(A) \setminus j(A) \cap E}$ is not birational onto its image and call $x \geq 2$ its degree. Thus $\deg(T) \leq d/x \leq d/2$. Since $j(A)$ spans $\mathbb{P}^r$, T spans $\mathbb{P}^{r-t}$. Hence $\deg(T) \geq r - t = d - p_a(A) - t$. Hence $(d - p_a(A) - t) \geq 2(d - p_a(A) - t)$, contradicting our assumption $d > p_a(A) + t$.

Lemma 2. Fix an integer $y > 0$. Let A be a geometrically integral projective curve defined over $\mathbb{F}_q$. Assume $A_{\text{reg}}(\mathbb{F}_q) \neq \emptyset$ and fix $P \in A_{\text{reg}}(\mathbb{F}_q)$. Then there are a

geometrically integral projective curve Y and a morphism $u : A \rightarrow Y$ defined over $\mathbb{F}_q$ such that u induces an isomorphism of $A \setminus \{P\}$ onto $Y \setminus u(P)$ and $p_a(Y) = p_a(A) + y$.

Proof. Let $\mathfrak{m}$ be the maximal ideal of the local ring $\mathcal{O}_{A,P}$. By assumption $\mathcal{O}_{A,P}/\mathfrak{m} \cong \mathbb{F}_q$ and $\mathbb{F}_q \cdot 1 \subset \mathcal{O}_{A,P}$. Hence the $\mathbb{F}_q$ -vector space $\mathcal{O}_{A,P}$ is the direct sum of its subspaces $\mathbb{F}_q \cdot 1$ and $\mathfrak{m}$. Set $\mathcal{O}_{Y,u(P)} := \mathbb{F}_q \cdot 1 + \mathfrak{m}^{y+1} \subset \mathcal{O}_{A,P}$. It is easy to check that $\mathcal{O}_{Y,u(P)}$ is a local ring with $\mathfrak{m}^{y+1}$ as its maximal ideal. Since $P \in A_{reg}$, $\mathcal{O}_{A,P}$ is a DVR. Hence $\mathfrak{m}/\mathfrak{m}^{t+1}$ is a $\mathbb{F}_q$ -vector space of dimension y . We take as Y the same topological space as Y , but with $\mathcal{O}_{Y,u(P)}$ at the point $u(P)$ associated to P instead of $\mathcal{O}_{A,P}$. With this definition of u we have $\dim_{\mathbb{F}_q}(u_*(\mathcal{O}_A)/\mathcal{O}_Y) = y$. Hence $p_a(Y) = p_a(A) + y$. $\square$

Remark 4. Fix q , C and an integer $n \geq 2$. Here we explain one way to check the existence of the curve $C_{[q,n]}$. We obtain $C_{[q,n]}$ in finitely many steps each of them similar to the one described in Example 2. We use z steps, where z is the number of orbits of F_q in $C(\mathbb{F}_{q^n}) \setminus C(\mathbb{F}_q)$. At each of the steps we glue together one of these orbit. We do not need any notion of gluing, except that set-theoretically in each step one of these orbits is sent to a single points and for all other points the map is an isomorphism. Fix $Q \in C(\mathbb{F}_{q^n}) \setminus C(\mathbb{F}_q)$ and assume $Q \in C(\mathbb{F}_{q^t}) \setminus C(\mathbb{F}_{q^{t-1}})$. Hence $\{Q, F_q(Q), \dots, F_q^{t-1}(Q)\}$ is the orbit of Q for the action of F_q . Call A the geometrically integral curve arising in the steps at which we want to glue this orbit. Hence there is a geometrically integral projective A curve defined over $\mathbb{F}_q$ with C as its normalization (call $u : C \rightarrow A$) and $u(Q) \in A_{reg}$ (in the previous steps (if any) the maps where isomorphism at each point of $\{Q, F_q(Q), \dots, F_q^{t-1}(Q)\}$). Set $P := u(Q)$. Since u is defined over $\mathbb{F}_q$ and u is an isomorphism in a neighborhood of $u^{-1}(\{P, F_q(P), \dots, F_q^{t-1}(P)\})$, we have $\{P, F_q(P), \dots, F_q^{t-1}(P)\} \subset A_{reg}$ and these t points are distinct. Hence $P \in A_{reg}(\mathbb{F}_{q^t}) \setminus A_{reg}(\mathbb{F}_{q^{t-1}})$. As in Example 2 we get several curves Y and morphism $v : A \rightarrow Y$ defined over $\mathbb{F}_q$, sending $\{P, F_q(P), \dots, F_q^{t-1}(P)\}$ to a single point, O , of Y and induces an isomorphism of $A \setminus \{P, F_q(P), \dots, F_q^{t-1}(P)\}$ onto $Y \setminus \{O\}$. Let A_1 be the seminormalization of Y in A . Then we use A_1 instead of A . After z steps we get $C_{[q,n]}$. To get $C_{[q,n]}$ we get an existence property for the seminormalization. The result does not depend from the order of the gluing. Hence $C_{[q,n]}$ depends only from q , C and n . Hence the curves $\mathbb{P}_{[q,n]}^1$ depends only from q and n .

REFERENCES

- [1] Y. Aubry and M. Perret, A Weil theorem for singular curves, in: R. Pellikaan, M. Perret and S. Vlăduț (Eds.), Arithmetic geometry and coding theory (Luminy, 1993), 1–7, de Gruyter, Berlin, 1996.
- [2] E. D. Davis, On the geometric interpretation of seminormality. Proc. Amer. Math. Soc. 68 (1978), no. 1, 1–5.
- [3] P. Deligne, La conjecture de Weil. II. Publ. Math. IHES 52 (1980), 137–252.
- [4] S. Fukasawa, Galois points for non-reflexive plane curve of low degree, preprint.
- [5] S. Fukasawa, M. Homma and S. J. Kim, Rational curves with many rational points over a finite field, arXiv:1108.4227.
- [6] Hirschfeld, J. W. P., Korchmáros and G. Torres, F.: Algebraic curves over a finite field. Princeton Series in Applied Mathematics. Princeton University Press, Princeton, NJ, 2008.
- [7] J.-P. Serre, Groupes algébriques et corps de classes. Hermann, Paris, 1959.
- [8] K.-O. Stöhr, On the poles of regular differentials of singular curves. Bol. Soc. Brasil. Mat. (N.S.) 24 (1993), no. 1, 105–136.
- [9] C. Traverso, Seminormality and Picard group. Ann. Scuola Norm. Sup. Pisa (3) 24 (1970), 585–595.

- [10] P. A. Viana and J. E. A. Rodrigues, Eventually minimal curves. Bull. Braz. Math. Soc. (N.S.) 36 (2005), no. 1, 39–58.
- [11] M. A. Vitulli, Weak normality and seminormality. Commutative algebra — Noetherian and non-Noetherian perspectives, 441–480, Springer, New York, 2011.
- [12] W. A. Zúñiga-Galindo, Number of rational points of a singular curve, Proc. Amer. Math. Soc. 126 (1998), no. 9, 2549–2556.

DEPT. OF MATHEMATICS, UNIVERSITY OF TRENTO, 38123 POVO (TN), ITALY
E-mail address: `ballico@science.unitn.it`